



# Measurements of the Higgs Boson at the LHC and Tevatron

Somnath Choudhury

(for the ATLAS, CMS, DØ and CDF collaborations)



44<sup>th</sup> International Symposium on Multiparticle Dynamics  
8 – 12 September 2014, Bologna (Italy)





Volume 712, Issue 3, 6 June 2012

ISSN 0370-2693

**5.1 fb<sup>-1</sup> @ 7 TeV + 5.3 fb<sup>-1</sup> @ 8 TeV**

# PHYSICS LETTERS B

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)  
SciVerse ScienceDirect





<http://www.elsevier.com/locate/physletb>


## Outline

- LHC, Tevatron & Detectors
- Higgs Observation in Boson and Fermion decay
- Higgs Combination
- Higgs Measurements – Mass, Coupling, Spin/Parity Width, Cross-section
- Summary - LHC, Tevatron

# The LHC



**CMS Integrated Luminosity, pp**




Overall data taking efficiency  $\sim 90\%$  by both detectors  
 Mean pileup 21 interactions/bunch crossing recorded  
 A major challenge overcome by both ATLAS and CMS

# The Tevatron





## The Tevatron accelerator



## Beam tunnel of Tevatron ring



**Proton-Antiproton collider Tevatron Run II:** (2002-2011),  $\sqrt{s} = 1.96$  TeV  
A decade of successful running delivered  $\sim 10 \text{ fb}^{-1}$  of luminosity for physics to CDF and DØ, shut down September 2011





## Detector subsystems

Vertex Detector  
 Silicon Tracker

Electromagnetic Calorimeter  
 Hadronic Calorimeter

Muon System

# DØ & CDF detector









# PHYSICS LETTERS B

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)





SciVerse ScienceDirect






# SM Higgs Sector @ LHC

Great achievement to four decade long quest  
A Higgs-like state pinned down at 125 GeV




**Higgs (125)**

# Higgs Observation








arXiv:1407.0558 [hep-ex]




- The signal @125 GeV with significance  $> 5\sigma$  observed in  $\gamma\gamma$  decay mode
- $6.7\sigma$  significance @125 GeV observed in  $H \rightarrow ZZ^* \rightarrow 4l$  golden decay mode



$$H \rightarrow \gamma\gamma$$





$$H \rightarrow ZZ^* \rightarrow 2e2\mu$$



## Observation of Higgs Boson in $\gamma\gamma$ and $ZZ^* \rightarrow 4\ell$ decay channels at ATLAS


arXiv:1408.7084 [hep-ex]

arXiv:1408.5191 [hep-ex]



- $H \rightarrow ZZ^* \rightarrow 4\ell$  signal observed with a significance of  $8.1\sigma$  at  $m_H = 125.36 \text{ GeV}$
- Measured signal strength  $\mu = 1.44^{+0.40}_{-0.33}$  in  $ZZ^* \rightarrow 4\ell$  and  $1.17 \pm 0.27$  in  $\gamma\gamma$  mode

$$H \rightarrow \gamma\gamma$$




 **ATLAS**  
EXPERIMENT

Run Number: 204769, Event Number: 24947130

Date: 2012-06-10 08:17:12 UTC

$$H \rightarrow ZZ^* \rightarrow 4\mu$$




Run Number: 204769  
Event Number: 82599793  
Date: 2012-06-10, 13:12:52 CET

EtCut > 0.4 GeV  
PtCut > 1.0 GeV


Muon: blue  
Cells: Tiles, EMC



Phys. Rev. Lett. 109, 071804 (2012)



Combined searches for  $WH \rightarrow l\nu b\bar{b}$ ,  $ZH \rightarrow ll b\bar{b}$ , and  $WH+ZH \rightarrow \text{MET} b\bar{b}$  using full  $9.7 \text{ fb}^{-1}$  Tevatron dataset




Excess of data over the background seen with most significance in the mass range  $120 \leq m_H \leq 135 \text{ GeV}$   
**1<sup>st</sup> Indication of Higgs-Fermion coupling**

**2.8 $\sigma$  significance at  $m_H = 125 \text{ GeV}$**

# Higgs - Lepton Coupling @ ATLAS


ATLAS-CONF-2013-108

## Evidence for Higgs - Lepton Coupling



- Excess  $> 4\sigma$  observed over  $m_H$  110 - 130 GeV
- Best fit signal strength  $1.4^{+0.5}_{-0.4}$  for  $m_H=125$  GeV
- Observed Significance  $4.1\sigma$  for  $m_H=125$  GeV
- A clear indication of Higgs - Lepton coupling


**ATLAS Prelim.**  
 $m_H = 125 \text{ GeV}$



$$H \rightarrow \tau\tau$$



Run: 214021  
Event: 269834309

2012-11-05 09:48:46 UTC



**ATLAS**  
EXPERIMENT

# Higgs - Lepton Coupling @ CMS



## Evidence for Higgs - Lepton Coupling


JHEP 05 (2014) 104

Excess  $>3\sigma$  observed over  $m_H$  110-130 GeV

H  $\rightarrow \tau\tau$  best fit signal strength  $0.78 \pm 0.27$  for  $m_H = 125$  GeV

Observed (expected) Significance  $3.2\sigma$  ( $3.7\sigma$ ) for  $m_H = 125$  GeV

Strong affirmation on Higgs-Fermion coupling, a clear indication of Higgs-Lepton coupling



Signal strength  $\mu$  compared to SM

$\mu = 0.78 \pm 0.27$


$H \rightarrow \tau\tau$




CMS Experiment at the LHC, CERN

Data recorded: 2012-Jun-05 09:58:43.400262 GMT(11:58:43 CEST)


Run / Event: 195552 / 61758463



Nature Physics 10, 557 (2014)



Channel ( $m_H = 125 \text{ GeV}$ )	Significance ( $\sigma$ )		Best-fit $\mu$
	Expected	Observed	
$VH \rightarrow b\bar{b}$	2.3	2.1	$1.0 \pm 0.5$
$H \rightarrow \tau\tau$	3.7	3.2	$0.78 \pm 0.27$
Combined	4.4	3.8	$0.83 \pm 0.24$





Strong evidence for the **direct coupling** of the **125 GeV Higgs boson** to **fermions**, with an **observed (expected) significance of  $3.8\sigma$  ( $4.4\sigma$ )**

- **ttH** important to probe directly top-Higgs Yukawa coupling
- Interested in ttH, H to anything with large enough BR  
H→bb is the most attractive here for SM  $m_H = 125$  GeV
- Categories events based on number of jets and b-tagged jets



arXiv:1408.1682 [hep-ex]

ATLAS-CONF-2014-043





# Higgs Combination





CMS-PAS-HIG-14-009

$$\sigma/\sigma_{\text{SM}} = 1.00 \pm 0.13 \left[ \pm 0.09(\text{stat.})_{-0.07}^{+0.08}(\text{theo.}) \pm 0.07(\text{syst.}) \right]$$




# Higgs Combination @ ATLAS

**ATLAS Prelim.**

$m_H = 125.5$  GeV

$\sigma(\text{stat.})$   
 $\sigma(\text{sys inc.})$   
 $\sigma(\text{theory})$

Total uncertainty  $\pm 1\sigma$  on  $\mu$




$\sqrt{s} = 7$  TeV  $\int \text{Ldt} = 4.6\text{-}4.8 \text{ fb}^{-1}$   
 $\sqrt{s} = 8$  TeV  $\int \text{Ldt} = 20.3 \text{ fb}^{-1}$

Signal strength ( $\mu$ )

ATLAS-CONF-2014-009

$\mu = 1.30 \pm 0.12$  (stat)  $^{+0.14}_{-0.11}$  (sys) @  $m_H = 125.5$  GeV





Observation in good agreement with the Standard Model Higgs boson hypothesis

# Higgs Combination @ Tevatron

Phys. Rev. D 88, 052014 (2013)



arXiv:1203.3774 [hep-ex]



- Combined searches by CDF and D0 for the SM Higgs boson in the mass range 100–200 GeV for  $bb$ ,  $\tau\tau$ ,  $\gamma\gamma$ ,  $ZZ$  and  $WW$  modes
- A significant excess of events observed in the mass range between 115 and 140 GeV with local significance at  $m_H = 125 \text{ GeV}$  corresponding to  $3.0\sigma$



# Higgs Mass







**H → γγ and H → ZZ\* → 4l**  
invariant mass distribution at CMS

# High Resolution Mass @ ATLAS



$H \rightarrow \gamma\gamma$  and  $H \rightarrow ZZ^* \rightarrow 4l$   
 invariant mass at ATLAS

# Higgs Mass



$H \rightarrow \gamma\gamma$  and  $H \rightarrow ZZ^* \rightarrow 4l$  combined mass measurement

$$m_H(\text{ATLAS}) = 125.36 \pm 0.37 \text{ (stat)} \pm 0.18 \text{ (syst)}$$

$$m_H(\text{CMS}) = 125.03 \begin{matrix} +0.26 \\ -0.27 \end{matrix} \text{ (stat)} \begin{matrix} +0.13 \\ -0.15 \end{matrix} \text{ (syst)}$$



# Higgs Coupling







CMS-PAS-HIG-14-009





2D likelihood scan for  $\kappa_g$  and  $\kappa_\gamma$  parameters assuming that  $\Gamma_{BSM}=0$



Results within  $1\sigma$  of Standard Model prediction

# Mass & Coupling with $H \rightarrow \tau\tau$



Scan of  $-2\Delta\ln L$ , as function of  $m_H$

background-only hypothesis includes the  $pp \rightarrow H(125 \text{ GeV}) \rightarrow WW$  process

**$M_H = 122 \pm 7 \text{ GeV}$**

Likelihood scan as a function of  $\kappa_V$  and  $\kappa_f$   
All nuisance param. profiled for each point

$pp \rightarrow H(125 \text{ GeV}) \rightarrow WW$  process added as a signal for vector boson coupling sensitivity

# Higgs Coupling @ ATLAS






ATLAS-CONF-2014-009

**ATLAS Preliminary**

$m_H = 125.5$  GeV

Total uncertainty


$\pm 1\sigma$  on  $\frac{\mu_{VBF+VH}}{\mu_{ggF+ttH}}$




# Higgs Coupling @ Tevatron



Phys. Rev. D 88, 052014 (2013)



The points that maximize the local posterior probability densities are marked with dots



2D constraints in  $(\kappa_W, \kappa_Z)$  plane for  $m_H=125 \text{ GeV}$  allowing  $\kappa_f$  to float

2D constraints in the  $(\kappa_V, \kappa_f)$  plane assuming Custodial symmetry ( $\lambda_{WZ}=1$ )





# Higgs Spin / Parity



## Matrix Element Likelihood Analysis




$$\text{MELA} = \left[ 1 + \frac{\mathcal{P}_{\text{bkg}}(m_1, m_2, \theta_1, \theta_2, \Phi, \theta^*, \Phi_1 | m_{4\ell})}{\mathcal{P}_{\text{sig}}(m_1, m_2, \theta_1, \theta_2, \Phi, \theta^*, \Phi_1 | m_{4\ell})} \right]^{-1}$$

Masses of dilepton pairs and five angles fully defining a four-lepton configuration in their centre-of-mass frame



# Higgs Spin & Parity @ CMS

## Discriminant for production and decay of different Higgs $J^P$ state



$$D_{J^P} = \frac{\mathcal{P}_{SM}}{\mathcal{P}_{SM} + \mathcal{P}_{J^P}} = \left[ 1 + \frac{\mathcal{P}_{J^P}(m_{Z_1}, m_{Z_2}, \vec{\Omega} | m_{4\ell})}{\mathcal{P}_{SM}(m_{Z_1}, m_{Z_2}, \vec{\Omega} | m_{4\ell})} \right]^{-1}$$

Statistically equivalent to the 2D analysis of  $m_{4\ell}$  and  $K_D$

$$D_{\text{bkg}} = \mathcal{P}_{\text{sig}} / (\mathcal{P}_{\text{sig}} + \dot{\mathcal{P}}_{\text{bkg}})$$


Hypotheses of a **pseudoscalar** and all tested spin-1 boson hypotheses excluded at 99% CL or higher

**Consistency with SM scalar boson**



CMS-PAS-HIG-14-014

- Using full angular information defining 4 lepton system
- For each hypothesis create kinematic discriminant for SM vs alternative hypothesis
- Perform 2D fit of hypothesis discriminant versus background discriminant and perform hypothesis test





Several J<sup>P</sup> hypotheses have been tested  
 Consistency with the SM scalar boson




# Higgs Spin & Parity @ ATLAS


Phys. Lett. B 726, 120 (2013)



Hypotheses of a **pseudoscalar** state as well as spin-1 and spin-2 hypotheses disfavoured over a SM scalar boson



# Higgs Width





Off-shell Higgs production sizeable at high ZZ mass

- ~7.6% of the total cross-section for  $m_{ZZ} > 2M_Z$
- Destructive interference between  $gg \rightarrow H \rightarrow ZZ$  and  $gg \rightarrow ZZ$

Ratio of **on-shell** and **off-shell** production cross-section gives a direct handle to constrain the total width (taking into account interference)

## On shell and off shell production in ZZ:

$$\sigma_{gg \rightarrow H \rightarrow ZZ}^{\text{on-shell}} \sim \frac{g_{ggH}^2 g_{HZZ}^2}{m_H \Gamma_H}$$

$$\sigma_{gg \rightarrow H \rightarrow ZZ}^{\text{off-shell}} \sim \frac{g_{ggH}^2 g_{HZZ}^2}{(2m_Z)^2}$$

On-shell and off-shell cross-section expressed as a function of signal strength by scaling the couplings

- **On-shell**

cross section constrained by  $H \rightarrow ZZ \rightarrow 4l$  search  
 $\mu = \sigma / \sigma_{SM} = 0.93^{+0.26}_{-0.24}$  (Expectation of  $1.0^{+0.27}_{-0.24}$ )

- **Off-shell**

cross section is constrained by  $H \rightarrow ZZ \rightarrow 4l$  and  $H \rightarrow ZZ \rightarrow 2l2\nu$  final states

Proposed by  
 F. Caola, K. Melnikov, PRD 88 (2013) 054024  
 N. Kauer, G. Passarino, JHEP 08 (2012) 116  
 J. Campbell et al. (arXiv:1311.3589)



## 4l final state

Using the baseline  $H \rightarrow ZZ$  selection strategy

- Exploiting full reconstructed final state to separate  $gg \rightarrow ZZ$  from  $qq \rightarrow ZZ$  at high mass
- Using angular discriminant as in the 4l baseline search
- Signal extracted by 2D fit in the mass and kinematic discriminant


## 2l2v final state

Requiring a di-lepton and high missing  $E_T$

- Mass shape fit in different jet categories

- **Observed limit of 5.4 x SM corresponding to ~ 22 MeV @ 95% CL**
- **Sensitivity exceeds all expectations enhancing the LHC role of a Higgs factory in next years**


Physics Letters B 736, 64 (2014)





# Higgs Width @ ATLAS

- Analysis in the 4l channel uses a likelihood fit to the distribution of a matrix element discriminant
- Analysis in the 2l2v channel counts events in a  $H^* \rightarrow ZZ$  enriched signal region with high  $E_T$  and high transverse mass
- Separate  $gg \rightarrow ZZ$  from  $qq \rightarrow ZZ$  at high mass

ATLAS-CONF-2014-042



$\Gamma/\Gamma_H^{\text{SM}}$  observed (expected) limit  
**4.8–7.7 (7.0–12.0) x SM @95% CL**



# Higgs Cross-section




arXiv:1407.4222 [hep-ex]

$H \rightarrow \gamma\gamma$



Measured cross sections and cross-section limits for  $pp \rightarrow H \rightarrow \gamma\gamma$  in seven fiducial regions

Data compared to state-of-the-art theoretical predictions, regions include SM prediction arising from VBF, VH and ttH labelled as XH




arXiv:1407.4222 [hep-ex]


Sensitive to new physics in the context of production loop and properties

**Jet multiplicity:** sensitive to relative rates of production modes (ggH, VBF/VH/ZH, ttH)

$\Delta\phi_{jj}$ : for ggH and VBF, sensitive to the Higgs boson spin and CP



$H \rightarrow \gamma\gamma$





- ❑ Higgs boson observed at the LHC and Tevatron
- ❑ Di-Tau Higgs decay at LHC- First Indication of Higgs coupling to Leptons
- ❑ Coupling and Spin/Parity properties measured show consistency with Standard Model
- ❑ Starting 2015, a robust Higgs Physics program

## CMS Projection

